



Autism Spectrum Disorder and Eating Problems: The Imbalance of Gut Microbiota and the Gut-Brain Axis Hypothesis

Jiyoung Kim^{1,2}

¹NUSEUM Inc., Seoul, Korea

²Center for Food and Bioconvergence, Seoul National University, Seoul, Korea

This review explores the complexities of autism spectrum disorder (ASD), primarily focusing on the significant eating challenges faced by children and adolescents with this neurodevelopmental condition. It is common for individuals with ASD to exhibit heightened sensitivity to various sensory aspects of food such as taste, texture, smell, and visual appeal, leading to restricted and less diverse diets. These dietary limitations are believed to contribute to an imbalance in the gut microbiota. This review elaborates on how these eating problems, coupled with the distinctive characteristics of ASD, might be influenced by and, in turn, influence the gut-brain axis, a bidirectional communication system between the gastrointestinal tract and the brain. This discussion aims to shed light on the multifaceted interactions and potential implications of diet, gut health, and neurological development and function in children and adolescents with ASD.

Keywords: Autism spectrum disorder; Eating disorder; Gastrointestinal tract; Microbiome; Gut-brain axis.

Received: October 8, 2023 / **Revised:** November 29, 2023 / **Accepted:** November 29, 2023

Address for correspondence: Jiyoung Kim, Center for Food and Bioconvergence, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Korea
Tel: +82-2-880-4921, Fax: +82-2-873-5260, E-mail: jiyoungkim1107@snu.ac.kr

INTRODUCTION

Autism spectrum disorder (ASD) is a developmental disorder marked by impairments in social interactions and communication skills, coupled with repetitive behaviors and restricted interests [1]. Many children and adolescents diagnosed with ASD experience significant eating challenges [2]. Many factors contribute to these dietary challenges, including enhanced sensitivity to the sensory qualities of food such as taste, texture, smell, and appearance [2]. Furthermore, individuals with ASD frequently find it difficult to incorporate new foods into their diet or adapt to various eating methods [2]. Many individuals find articulating food preferences or dietary needs daunting, which commonly leads to anxiety related to mealtimes [2]. Children with ASD who display restricted interests tend to consume limited diets [3] indicating that restricted interest could induce an imbalance in the gut microbiota of children and adolescents with ASD.

ASD AND FOOD SELECTIVITY

Food selectivity is a prevalent eating challenge in ASD, ex-

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

posing children to heightened risks of inadequate intake of vital minerals and vitamins, which impacts their developmental trajectory [4]. Furthermore, pronounced food selectivity in toddlers with ASD correlates with the intensification of autism symptoms [4].

Sensory sensitivity

Atypical sensory processing in ASD represents a crucial, albeit less frequently discussed, subject of discourse. This phenomenon is conspicuous across various sensory realms, including touch, smell, and taste, and manifests with greater prevalence in children with ASD than in those exhibiting other developmental disabilities or those with typical development [5]. The resultant sensory sensitivities often materialize as under-responsivity (diminished reactions to sensory stimuli), over-responsivity (amplified or adverse reactions to sensory stimuli), or a composite of both. These sensory idiosyncrasies are enduring, unaffected by age or ASD severity, can be discernible from early childhood, and persist throughout the individual's lifetime [5].

Sensitivity, specifically within the oral sensory domain, has pronounced implications for dietary behaviors. There is a link between food selectivity in ASD and sensory processing malfunctions with a specific focus on oral sensory sensitivity [4-8]. Children with distinctive oral sensory sensitivi-

ties exhibit broader rejection of food varieties and diminished intake of vegetables than those devoid of such sensitivities [9]. Conversely, a deficiency in taste sensitivity is associated with increased symptoms of eating disorders [4]. Moreover, enhanced sensitivity in the visual domain correlates with heightened symptoms of both eating disorders and ASD-specific eating habits [4].

An in-depth investigation into sensory sensitivities and their diverse implications for the dietary patterns of children and adolescents with ASD is vital, given the intricate relationship between sensory processing and dietary choices. A comprehensive understanding of their varied manifestations and the formulation of customized intervention strategies designed to accommodate the unique requirements of those navigating the complexities of the autism spectrum is required to effectively address these sensory peculiarities.

Neophobia

Food selectivity and neophobia manifest in children with ASD due to sensory sensitivity, adaptability issues, and anxiety [9]. A comprehensive review titled “*Food Selectivity and Neophobia in Children with Autism Spectrum Disorder and Neurotypical Development*” demonstrated that children with ASD are more predisposed to exhibit food selectivity and neophobia than their neurotypically developing peers [9]. This study highlighted the considerable impact of such behaviors on the health and well-being of children with ASD, indicating risks such as malnutrition, being overweight, social isolation, and difficulties in social participation [9]. Due to the substantial implications for the health, development, and social inclusion of individuals within the spectrum, the manifestations of food selectivity and neophobia in ASD require meticulous consideration. Recognizing the multifaceted roots of these phenomena is pivotal for formulating effective and holistic interventions tailored to the unique needs and challenges faced by children with ASD.

Restrictive behavior

Restrictive behavior is another common characteristic of ASD that can significantly affect food choices [4]. This behavior may manifest as a rigid adherence to specific routines or rituals, leading to a limited range of acceptable foods and a refusal to try new ones. Furthermore, children with ASD may exhibit avoidant/restrictive food intake disorder (ARFID), a newly recognized eating disorder characterized by a persistent pattern of food restriction that is often unrelated to other medical conditions or cultural norms [10]. ARFID can lead to severe nutritional deficiencies, dependence on oral supplements, and the need for enteral feeding [10].

The precise definition of “food selectivity” in the context of

ASD remains a subject of debate. However, studies suggest that rigid adherence to routines and rituals, along with sensory sensitivity and neophobia, contributes to the development of food selectivity in patients with ASD. A comprehensive approach is required to address these restrictive behaviors, considering their underlying causes and implementing individualized interventions.

DIET QUALITY AND NUTRITIONAL STATUS

Preschoolers with ASD often encounter significant challenges during mealtime, hindering their intake of essential nutrients and increasing their risk of nutritional deficiencies [11]. A case-control study comparing mealtime behaviors between Chinese preschoolers with ASD and typically developing preschoolers in Hong Kong revealed that the ASD group exhibited a significantly higher likelihood of food refusal and unhealthy eating habits [11]. This was evident in their higher scores on the Brief Autism Mealtime Behavior Inventory food refusal domain and the Chinese Preschoolers’ Eating Behavior Questionnaire food fussiness and eating habit domains [11]. Additionally, the ASD group was less responsive to external factors that influenced eating and was less inclined to eat independently [11].

There is a pervasive trend among children and adolescents with ASD to maintain a diet of inferior quality compared with national standard datasets as shown by a cross-sectional study [12]. This study highlighted notable disparities in diet quality, with ASD individuals exhibiting higher consumption of whole grains, fatty acids, added sugars, and refined grains and a lower intake of whole fruits, total vegetables, dairy, total protein foods, seafood, and plant proteins [12].

An extensive meta-analysis of the dietary patterns of children with ASD inferred that this population had a lower intake of proteins, calcium, phosphorus, selenium, vitamin D, thiamine, riboflavin, and vitamin B12 than typically developing children [13,14]. In contrast, they consumed an elevated level of polyunsaturated fatty acids and vitamin E [13,14]. Children with ASD also have lower omega-3 consumption than those without, necessitating cautious interpretation of these results because of the limited number of studies included in the analysis and high heterogeneity [13,14]. The findings also indicate deviations from the Dietary Reference Intake recommendations, with lower intakes of calcium, vitamin D, and dairy and higher intakes of protein, phosphorus, selenium, thiamine, riboflavin, and vitamin B12 than recommended [13,14].

There have been two reported cases of children with ASD becoming critically ill because of vitamin deficiencies [15].

In the first instance, the child suffered from thiamine deficiency, culminating in beriberi, which can induce heart complications, nerve damage, and other severe health issues [15]. Despite prompt vitamin supplementation, the patient experienced a cardiac arrest and developed serious neurological complications [15]. Another child displayed symptoms of scurvy resulting from vitamin C deficiency, manifesting as leg pain and a limping gait [15]. In this case, the administration of intravenous vitamin C and tube feeding led to a quick recovery [15]. The authors emphasized the existence of other reports of critically ill children with ASD due to thiamine deficiency and underscored the necessity of early interventions to prevent severe vitamin deficiencies arising from the observed high food selectivity in children with ASD [15]. This is a clear demonstration of the link between food selectivity and vitamin deficiency in children with ASD [15]. Case reports have provided compelling evidence supporting the hypothesis that selective eating in individuals with ASD can lead to vitamin deficiency [15]. The detailed descriptions of children's diets and the subsequent development of vitamin deficiencies underscore the importance of addressing selective eating behaviors in ASD to ensure adequate nutrition and overall health.

Children with ASD often exhibit lower vitamin D levels at birth than their non-ASD siblings likely due to a combination of genetic, environmental, and dietary factors and potentially influencing ASD development [16]. Therefore, lower vitamin D levels may be correlated with the severity of autism symptoms, emphasizing the imperative need for extensive research to clarify the relationship between vitamin D levels and ASD and develop novel treatments focusing on vitamin D levels [16].

ASD AND MICROBIOTA-GUT-BRAIN AXIS HYPOTHESIS

The gut maintains multifaceted interactions with the brain via diverse pathways. It utilizes neuroactive molecules and microbial metabolites, such as short-chain fatty acids and neurotransmitters produced by entities such as enteroendocrine cells, as chemical intermediaries [1]. These substances can modulate the central nervous system activity either directly by traversing the blood-brain barrier or indirectly by influencing components such as the vagus nerve and the mucosal immune response [1]. The gut microbiota, including *Bifidobacterium* and *Prevotella*, maintain chemical dialogues with the brain through compounds such as short-chain fatty acids and serotonin (5-hydroxytryptamine), which significantly affect mood, behavior, and cognition by modulating the production of neurotransmitters and interacting with var-

ious physiological components [1].

Emerging evidence suggests that the microbiota-gut-brain axis plays a crucial role in the development and progression of ASD, with children with ASD exhibiting variations in gut microbiota composition [3,17-23]. Alterations in the microbiota-gut-brain axis lead to dysbiosis of the gut microbiota, resulting in an imbalance in chemical signals and cytokines, which, in turn, heightens neuroinflammation in the brain in individuals with ASD [1]. Microglial activation and neurochemical dysregulation, including anomalies in glutamate, gamma-aminobutyric acid, serotonin, and oxytocin levels in the ASD brain, are correlated with disturbances in the gut microbiota [1]. A functional architecture along the gut-brain axis correlates with the heterogeneity of ASD phenotypes [24]. This architecture is characterized by ASD-associated amino acid, carbohydrate, and lipid profiles, which are predominantly encoded by microbial species belonging to the genera *Prevotella*, *Bifidobacterium*, *Desulfovibrio*, and *Bacteroides* [24]. Furthermore, this architecture correlates with changes in brain gene expression, restrictive dietary patterns, and proinflammatory cytokine profiles [24]. A strong association has been observed between temporal changes in the microbiome composition and ASD phenotypes [24].

There are associations between the intake of certain foods and the presence of certain bacteria in the gut microbiota [21]. For example, the consumption of dairy products is associated with higher levels of *Lactobacillus*, whereas the intake of cereals is associated with lower levels of *Roseburia* [21]. Children with ASD who are overweight or obese have been found to have lower levels of *Roseburia* and *Faecalibacterium prausnitzii* in their gut microbiota [21]. These bacteria produce beneficial fatty acids and other metabolites [21]. Therefore, dietary interventions may improve the gut and overall health of children with ASD. However, further research is required to develop and test specific dietary interventions for patients with ASD. Importantly, diet is just one of many elements that may influence the gut microbiome. Other factors, such as genetics, environment, and medications, may also play a role.

Specific instances of altered gut microbiota in ASD include a diminished presence of butyrate-producing bacteria such as *Ruminococcaceae*, *Eubacterium*, *Lachnospiraceae*, and *Erysipelotrichaceae* and decreased levels of gamma-aminobutyric acid-synthesizing microbiomes such as *Bacteroides*, *Lactobacillus*, and *Bifidobacterium* [1]. Such alterations highlight the promise of therapeutic interventions, such as probiotics and dietary adjustments, to improve the gut microbiota and consequently alleviate ASD symptoms.

Table 1. List of literatures covering ASD and eating problems

Eating problem	[2, 11, 25-30]
Sensory sensitivity	[5, 6, 27, 31-33]
Food selectivity	[4-6, 9, 10, 30, 34-55]
Nutritional intake	[13, 14, 21, 56-62]
Diet quality	[11, 12, 29, 63, 64]
Nutritional status	[21, 57-60, 65-76]
Others	[2, 3, 6, 11, 15, 16, 19, 28, 42, 57, 61, 63, 66, 75, 77-86]

ASD, autism spectrum disorder

CONCLUSION

Despite the extensive research and literature on ASD and eating problems (Table 1), there remains a need for greater attention to the management of these conditions.

There is a multifaceted relationship between specific gut microbiota, the gut-brain axis, and ASD, suggesting that improving gut health can be pivotal in alleviating ASD symptoms, with dietary interventions standing out as promising therapeutic strategies. Progressive research has elucidated the intricate pathways and significant roles of specific gut microbiota and the brain in regulating multiple brain functions, offering potential advancements in the understanding and management of ASD and its associated conditions, thus reinforcing the significance of continuous exploration in this complex biological domain.

Availability of Data and Material

Data sharing is not applicable to this article, as no datasets were generated or analyzed during the study.

Conflicts of Interest

The author has no potential conflicts of interest to disclose.

ORCID ID

Jiyoung Kim <https://orcid.org/0000-0002-2999-3320>

Funding Statement

None

Acknowledgments

I would like to express my sincere gratitude to Amazon Web Service Korea, MinSung Cho, and Sungwon Yang for their support in developing innovative digital technologies to address eating problems among children with ASD.

REFERENCES

- 1) Yu Y, Zhao F. Microbiota-gut-brain axis in autism spectrum disorder. *J Genet Genomics* 2021;48:755-762.
- 2) Baraskewich J, von Ranson KM, McCrimmon A, McMorris CA. Feeding and eating problems in children and adolescents with autism: a scoping review. *Autism* 2021;25:1505-1519.
- 3) Yap CX, Henders AK, Alvares GA, Wood DLA, Krause L, Tyson GW, et al. Autism-related dietary preferences mediate autism-gut microbiome associations. *Cell* 2021;184:5916-5931.e17.
- 4) Stolar O, Zachor DA, Ben-Itzhak E. Food selectivity is associated with more severe autism symptoms in toddlers with autism spectrum disorder. *Acta Paediatr* 2021;110:3305-3307.
- 5) Chistol LT, Bandini LG, Must A, Phillips S, Cermak SA, Curtin C. Sensory sensitivity and food selectivity in children with autism spectrum disorder. *J Autism Dev Disord* 2018;48:583-591.
- 6) Cermak SA, Curtin C, Bandini LG. Food selectivity and sensory sensitivity in children with autism spectrum disorders. *J Am Diet Assoc* 2010;110:238-246.
- 7) Suarez MA. Sensory processing in children with autism spectrum disorders and impact on functioning. *Pediatr Clin North Am* 2012;59:203-214.
- 8) Zobel-Lachiusa J, Andrianopoulos MV, Mailloux Z, Cermak SA. Sensory differences and mealtime behavior in children with autism. *Am J Occup Ther* 2015;69:6905185050.
- 9) Rodrigues JVS, Poli MCF, Petrilli PH, Dornelles RCM, Turcio KH, Theodoro LH. Food selectivity and neophobia in children with autism spectrum disorder and neurotypical development: a systematic review. *Nutr Rev* 2023;81:1034-1050.
- 10) Bourne L, Mandy W, Bryant-Waugh R. Avoidant/restrictive food intake disorder and severe food selectivity in children and young people with autism: a scoping review. *Dev Med Child Neurol* 2022;64:691-700.
- 11) Yeung SSY, Chan R, Li L, Chan D, Leung J, Leung TF. Eating behaviors and diet quality in Chinese preschoolers with and without autism spectrum disorder: a case-control study. *J Pediatr* 2021;237:258-266.e5.
- 12) Buro AW, Gray HL, Kirby RS, Berkman K, Agazzi H, Shaffer-Hudkins E. Diet quality in an ethnically diverse sample of children and adolescents with autism spectrum disorder compared with nationally representative data. *Disabil Health J* 2021;14:100981.
- 13) Esteban-Figuerola P, Canals J, Fernández-Cao JC, Arija Val V. Corrigendum to differences in food consumption and nutritional intake between children with autism spectrum disorders and typically developing children: a meta-analysis. *Autism* 2020;24:531-536.
- 14) Esteban-Figuerola P, Canals J, Fernández-Cao JC, Arija Val V. Differences in food consumption and nutritional intake between children with autism spectrum disorders and typically developing children: a meta-analysis. *Autism* 2019;23:1079-1095.
- 15) Chong PF, Torio M, Fujii F, Hirata Y, Matsuoka W, Sonoda Y, et al. Critical vitamin deficiencies in autism spectrum disorder: reversible and irreversible outcomes. *Eur J Clin Nutr* 2022;76:1618-1621.
- 16) Fernell E, Bejerot S, Westerlund J, Miniscalco C, Simila H, Eyles D, et al. Autism spectrum disorder and low vitamin D at birth: a sibling control study. *Mol Autism* 2015;6:3.
- 17) Mangiola F, Ianiro G, Franceschi F, Fagioli S, Gasbarrini G, Gasbarrini A. Gut microbiota in autism and mood disorders. *World J Gastroenterol* 2016;22:361-368.
- 18) Barnhill K, Gutierrez A, Ghossainy M, Marediya Z, Marti CN, Hewitson L. Growth status of children with autism spectrum disorder: a case-control study. *J Hum Nutr Diet* 2017;30:59-65.
- 19) Sanlier N, Kocabas S. The effect of probiotic, prebiotic and gut microbiota on ASD: a review and future perspectives. *Crit Rev Food Sci Nutr* 2023;63:2319-2330.
- 20) Alshammari MK, AlKhulaifi MM, Al Farraj DA, Somily AM, Al-Barrag AM. Incidence of *Clostridium perfringens* and its toxin genes in the gut of children with autism spectrum disorder. *Anaerobe* 2020;61:102114.
- 21) Mendive Dubourdieu P, Guendiain M. Understanding the link between gut microbiota, dietary intake, and nutritional status in children with autism and typical development. *Front Nutr* 2023;10:1202948.
- 22) Wan Y, Zuo T, Xu Z, Zhang F, Zhan H, Chan D, et al. Underdevel-

- opment of the gut microbiota and bacteria species as non-invasive markers of prediction in children with autism spectrum disorder. *Gut* 2022;71:910-918.
- 23) **Larroya-García A, Navas-Carrillo D, Orenes-Piñero E.** Impact of gut microbiota on neurological diseases: diet composition and novel treatments. *Crit Rev Food Sci Nutr* 2019;59:3102-3116.
 - 24) **Morton JT, Jin DM, Mills RH, Shao Y, Rahman G, McDonald D, et al.** Multi-level analysis of the gut-brain axis shows autism spectrum disorder-associated molecular and microbial profiles. *Nat Neurosci* 2023;26:1208-1217.
 - 25) **Keown K, Bothwell J, Jain S.** Nutritional implications of selective eating in a child with autism spectrum disorder. *BMJ Case Rep* 2014;2014:bcr2013202581.
 - 26) **Kozak A, Czepczor-Bernat K, Modrzejewska J, Modrzejewska A, Matusik E, Matusik P.** Avoidant/restrictive food disorder (ARFID), food neophobia, other eating-related behaviours and feeding practices among children with autism spectrum disorder and in non-clinical sample: a preliminary study. *Int J Environ Res Public Health* 2023;20:5822.
 - 27) **Nisticò V, Faggioli R, Tedesco R, Giordano B, Priori A, Gambini O, et al.** Brief report: sensory sensitivity is associated with disturbed eating in adults with autism spectrum disorders without intellectual disabilities. *J Autism Dev Disord* 2023;53:3295-3300.
 - 28) **Schreck KA, Williams K, Smith AF.** A comparison of eating behaviors between children with and without autism. *J Autism Dev Disord* 2004;34:433-438.
 - 29) **Kral TV, Eriksen WT, Souders MC, Pinto-Martin JA.** Eating behaviors, diet quality, and gastrointestinal symptoms in children with autism spectrum disorders: a brief review. *J Pediatr Nurs* 2013;28:548-556.
 - 30) **Floríndez LI, Floríndez DC, Price ME, Floríndez FM, Como DH, Polido JC, et al.** Exploring eating challenges and food selectivity for Latinx children with and without autism spectrum disorder using qualitative visual methodology: implications for oral health. *Int J Environ Res Public Health* 2021;18:3751.
 - 31) **Corbett BA, Schupp CW, Levine S, Mendoza S.** Comparing cortisol, stress, and sensory sensitivity in children with autism. *Autism Res* 2009;2:39-49.
 - 32) **Favre MR, Markram H, Markram K.** Individual differences in sensory sensitivity: further lessons from an autism model. *Cogn Neurosci* 2019;10:171-173.
 - 33) **Sapey-Triomphe LA, Leiros Costa T, Wagemans J.** Sensory sensitivity in autism mostly depends on contextual predictions. *Cogn Neurosci* 2019;10:162-164.
 - 34) **Schreck KA, Williams K.** Food preferences and factors influencing food selectivity for children with autism spectrum disorders. *Res Dev Disabil* 2006;27:353-363.
 - 35) **Bandini LG, Anderson SE, Curtin C, Cermak S, Evans EW, Scampini R, et al.** Food selectivity in children with autism spectrum disorders and typically developing children. *J Pediatr* 2010;157:259-264.
 - 36) **Mari-Bauset S, Zazpe I, Mari-Sanchis A, Llopis-González A, Morales-Suárez-Varela M.** Food selectivity in autism spectrum disorders: a systematic review. *J Child Neurol* 2014;29:1554-1561.
 - 37) **Suarez MA, Nelson NW, Curtis AB.** Longitudinal follow-up of factors associated with food selectivity in children with autism spectrum disorders. *Autism* 2014;18:924-932.
 - 38) **Curtin C, Hubbard K, Anderson SE, Mick E, Must A, Bandini LG.** Food selectivity, mealtime behavior problems, spousal stress, and family food choices in children with and without autism spectrum disorder. *J Autism Dev Disord* 2015;45:3308-3315.
 - 39) **Kuschner ES, Eisenberg IW, Orionzi B, Simmons WK, Kenworthy L, Martin A, et al.** A preliminary study of self-reported food selectivity in adolescents and young adults with autism spectrum disorder. *Res Autism Spectr Disord* 2015;15-16:53-59.
 - 40) **Postorino V, Sanges V, Giovagnoli G, Fatta LM, De Peppo L, Ar-mando M, et al.** Clinical differences in children with autism spectrum disorder with and without food selectivity. *Appetite* 2015;92:126-132.
 - 41) **Ma NS, Thompson C, Weston S.** Brief report: scurvy as a manifestation of food selectivity in children with autism. *J Autism Dev Disord* 2016;46:1464-1470.
 - 42) **Bandini LG, Curtin C, Phillips S, Anderson SE, Maslin M, Must A.** Changes in food selectivity in children with autism spectrum disorder. *J Autism Dev Disord* 2017;47:439-446.
 - 43) **Sharp WG, Postorino V, McCracken CE, Berry RC, Criado KK, Burrell TL, et al.** Dietary intake, nutrient status, and growth parameters in children with autism spectrum disorder and severe food selectivity: an electronic medical record review. *J Acad Nutr Diet* 2018;118:1943-1950.
 - 44) **Accardo J.** Food selectivity in autism: expanding the palate (and palette). *J Pediatr* 2019;211:1-3.
 - 45) **Hillman H.** Home-based video modeling on food selectivity of children with an autism spectrum disorder. *Phys Occup Ther Pediatr* 2019;39:629-641.
 - 46) **Smile S.** A 4-year-old boy with food selectivity and autism-spectrum disorder. *CMAJ* 2019;191:E636-E637.
 - 47) **Molina-López J, Leiva-García B, Planells E, Planells P.** Food selectivity, nutritional inadequacies, and mealtime behavioral problems in children with autism spectrum disorder compared to neurotypical children. *Int J Eat Disord* 2021;54:2155-2166.
 - 48) **Reche-Olmedo L, Torres-Collado L, Compañ-Gabucio LM, Garcia-de-la-Hera M.** The role of occupational therapy in managing food selectivity of children with autism spectrum disorder: a scoping review. *Children (Basel)* 2021;8:1024.
 - 49) **Pubyliki-Yanofchick W, Zaki-Scarpa C, LaRue RH, Manente C, Kahng S.** Treatment of food selectivity in an adult with autism spectrum disorder. *Behav Anal Pract* 2022;15:796-803.
 - 50) **Valenzuela-Zamora AF, Ramírez-Valenzuela DG, Ramos-Jiménez A.** Food selectivity and its implications associated with gastrointestinal disorders in children with autism spectrum disorders. *Nutrients* 2022;14:2660.
 - 51) **Beighley JS, Matson JL, Rieske RD, Adams HL.** Retraction notice to “food selectivity in children with and without an autism spectrum disorder: investigation of diagnosis and age” [Research in Developmental Disabilities 34/10 (2013) 3497-3503]. *Res Dev Disabil* 2023;136:104449. Retracted on May 2023. doi: 10.1016/j.ridd.2023.104449.
 - 52) **Bresciani G, Da Lozzo P, Lega S, Bramuzzo M, Di Leo G, Dissegna A, et al.** Gastrointestinal disorders and food selectivity: relationship with sleep and challenging behavior in children with autism spectrum disorder. *Children (Basel)* 2023;10:253.
 - 53) **Byrska A, Błażejczyk I, Faruga A, Potaczek M, Wilczyński KM, Janas-Kozik M.** Patterns of food selectivity among children with autism spectrum disorder. *J Clin Med* 2023;12:5469.
 - 54) **Espósito M, Mirizzi P, Fadda R, Pirollo C, Ricciardi O, Mazza M, et al.** Food selectivity in children with autism: guidelines for assessment and clinical interventions. *Int J Environ Res Public Health* 2023;20:5092.
 - 55) **Prosperi M, Santocchi E, Balboni G, Narzisi A, Bozza M, Fulceri F, et al.** Behavioral phenotype of ASD preschoolers with gastrointestinal symptoms or food selectivity. *J Autism Dev Disord* 2017;47:3574-3588.
 - 56) **Herndon AC, DiGuseppi C, Johnson SL, Leiferman J, Reynolds A.** Does nutritional intake differ between children with autism spectrum disorders and children with typical development? *J Autism Dev Disord* 2009;39:212-222.
 - 57) **Xia W, Zhou Y, Sun C, Wang J, Wu L.** A preliminary study on nutritional status and intake in Chinese children with autism. *Eur J Pediatr* 2010;169:1201-1206.
 - 58) **Padmanabhan PS, Shroff H.** The relationship between sensory in-

- tegration challenges and the dietary intake and nutritional status of children with autism spectrum disorders in Mumbai, India. *Int J Dev Disabil* 2018;66:142-152.
- 59) **Mendive Dubourdieu P, Guerendiain M.** Dietary intake, nutritional status and sensory profile in children with autism spectrum disorder and typical development. *Nutrients* 2022;14:2155.
 - 60) **Hernández Fernández IG, Omaña Covarrubias A, Conde Vega J, Rodríguez Ávila J, Díaz Martínez G, Moya Escalera A, et al.** [Nutritional status and food intake frequency in children with autism spectrum disorder]. *Nutr Hosp* 2023;40:347-353. Spanish
 - 61) **Sharp WG, Berry RC, McCracken C, Nuhu NN, Marvel E, Saulnier CA, et al.** Feeding problems and nutrient intake in children with autism spectrum disorders: a meta-analysis and comprehensive review of the literature. *J Autism Dev Disord* 2013;43:2159-2173.
 - 62) **Hyman SL, Stewart PA, Schmidt B, Cain U, Lemcke N, Foley JT, et al.** Nutrient intake from food in children with autism. *Pediatrics* 2012;130(Suppl 2):S145-S153.
 - 63) **Johnson CR, Turner K, Stewart PA, Schmidt B, Shui A, Macklin E, et al.** Relationships between feeding problems, behavioral characteristics and nutritional quality in children with ASD. *J Autism Dev Disord* 2014;44:2175-2184.
 - 64) **Li XS, Pinto-Martin JA, Thompson A, Chittams J, Kral TVE.** Weight status, diet quality, perceived stress, and functional health of caregivers of children with autism spectrum disorder. *J Spec Pediatr Nurs* 2018;23:e12205.
 - 65) **Zimmer MH, Hart LC, Manning-Courtney P, Murray DS, Bing NM, Summer S.** Food variety as a predictor of nutritional status among children with autism. *J Autism Dev Disord* 2012;42:549-556.
 - 66) **Sun C, Xia W, Zhao Y, Li N, Zhao D, Wu L.** Nutritional status survey of children with autism and typically developing children aged 4-6 years in Heilongjiang Province, China. *J Nutr Sci* 2013;2:e16.
 - 67) **Goldschmidt J.** Nutritional status and autism spectrum disorders. *Adv Nutr* 2015;6:865.
 - 68) **Mari-Bauset S, Llopis-González A, Zazpe-García I, Mari-Sanchis A, Morales-Suárez-Varela M.** Nutritional status of children with autism spectrum disorders (ASDs): a case-control study. *J Autism Dev Disord* 2015;45:203-212.
 - 69) **Ranjan S, Nasser JA.** Nutritional status of individuals with autism spectrum disorders: do we know enough? *Adv Nutr* 2015;6:397-407.
 - 70) **Mari-Bauset S, Llopis-González A, Zazpe I, Mari-Sanchis A, Morales Suárez-Varela M.** Comparison of nutritional status between children with autism spectrum disorder and typically developing children in the Mediterranean Region (Valencia, Spain). *Autism* 2017;21:310-322.
 - 71) **Altamimi M.** Could autism be associated with nutritional status in the Palestinian population? The outcomes of the Palestinian micro-nutrient survey. *Nutr Metab Insights* 2018;11:1178638818773078.
 - 72) **Zhu J, Guo M, Yang T, Lai X, Tang T, Chen J, et al.** Nutritional status and symptoms in preschool children with autism spectrum disorder: a two-center comparative study in Chongqing and Hainan province, China. *Front Pediatr* 2020;8:469.
 - 73) **Alkhalidy H, Abushaikha A, Alnaser K, Obeidat MD, Al-Shami I.** Nutritional status of pre-school children and determinant factors of autism: a case-control study. *Front Nutr* 2021;8:627011.
 - 74) **Lindsay RL, Eugene Arnold L, Aman MG, Vitiello B, Posey DJ, McDougle CJ, et al.** Dietary status and impact of risperidone on nutritional balance in children with autism: a pilot study. *J Intellect Dev Disabil* 2006;31:204-209.
 - 75) **Adams JB, Audhya T, McDonough-Means S, Rubin RA, Quig D, Geis E, et al.** Nutritional and metabolic status of children with autism vs. neurotypical children, and the association with autism severity. *Nutr Metab (Lond)* 2011;8:34.
 - 76) **Kawicka A, Regulska-Ilow B.** How nutritional status, diet and dietary supplements can affect autism. A review. *Rocz Panstw Zakl Hig* 2013;64:1-12.
 - 77) **Hubbard KL, Anderson SE, Curtin C, Must A, Bandini LG.** A comparison of food refusal related to characteristics of food in children with autism spectrum disorder and typically developing children. *J Acad Nutr Diet* 2014;114:1981-1987.
 - 78) **Emond A, Emmett P, Steer C, Golding J.** Feeding symptoms, dietary patterns, and growth in young children with autism spectrum disorders. *Pediatrics* 2010;126:e337-e342.
 - 79) **Sharp WG, Jaquess DL, Lukens CT.** Multi-method assessment of feeding problems among children with autism spectrum disorders. *Res Autism Spectr Disord* 2013;7:56-65.
 - 80) **Park JY, Kim YR, Lee SH, Kim Y, Oh JE, Chung SJ, et al.** A literature review of eating behaviors and nutrient intake of children with autism spectrum disorders. *J Korean Assoc Persons Autism* 2019;19:111-132.
 - 81) **Altun H, Kurutaş EB, Şahin N, Güngör O, Fındıklı E.** The levels of vitamin D, vitamin D receptor, homocysteine and complex B vitamin in children with autism spectrum disorders. *Clin Psychopharmacol Neurosci* 2018;16:383-390.
 - 82) **Mostafa GA, Al-Ayadhi LY.** Reduced serum concentrations of 25-hydroxy vitamin D in children with autism: relation to autoimmunity. *J Neuroinflammation* 2012;9:201.
 - 83) **Kałużna-Czaplińska J, Jóźwik-Pruska J.** Nutritional strategies and personalized diet in autism-choice or necessity? *Trends Food Sci Technol* 2016;49:45-50.
 - 84) **Lord C, Charman T, Havdahl A, Carbone P, Anagnostou E, Boyd B, et al.** The Lancet Commission on the future of care and clinical research in autism. *Lancet* 2022;399:271-334.
 - 85) **Weir E, Allison C, Ong KK, Baron-Cohen S.** An investigation of the diet, exercise, sleep, BMI, and health outcomes of autistic adults. *Mol Autism* 2021;12:31.
 - 86) **Adams JB, George F, Audhya T.** Abnormally high plasma levels of vitamin B6 in children with autism not taking supplements compared to controls not taking supplements. *J Altern Complement Med* 2006;12:59-63.